



# Coordinated Science Laboratory



UNIVERSITY OF ILLINOIS - URBANA, ILLINOIS

# **PSEUDO CUTS AND THEIR APPLICATIONS**

**Wataru Mayeda**

**REPORT R-233**

**JULY, 1964**

**COORDINATED SCIENCE LABORATORY  
UNIVERSITY OF ILLINOIS  
URBANA, ILLINOIS**

The research reported in this document was made possible by support extended to the University of Illinois, Coordinated Science Laboratory, under the Joint Services Electronics Program by the Department of the Army, Department of the Navy (Office of Naval Research), and the Department of the Air Force (Office of Scientific Research), and by the Advanced Research Projects Agency under Department of the Army contract

DA-28-043-AMC-00073(E)

and Department of the Air Force (Office of Scientific Research) contract

AF 49(638) - 1383

PSUEDO CUTS AND THEIR APPLICATION

Wataru Mayeda



## INTRODUCTION

Consider a non-oriented planar graph  $G$  and the dual graph  $[1,2,3]$   $G'$  of  $G$ . It is known that a circuit of  $G$  is a cut set of  $G'$ . Here we will introduce a set called a "psuedo cut" so that a special psuedo cut in  $G'$  is a path in  $G$ . When a linear graph is non-planar, a psuedo cut will have no relationship with a path in a linear graph. However, there is a similarity between the properties of paths and psuedo cuts. Let  $C$  be a class of all possible circuits, edge disjoint unions of circuits in a linear graph  $G$  and the empty set  $\emptyset$ . Let  $p$  be a path between vertices  $t$  and  $u$  in  $G$ . We define  $C\Phi_{tu}$  as

$$C\Phi_{tu} = \{x: x = p \oplus c_r, c_r \in C\}$$

Then all possible paths between  $t$  and  $u$  in  $G$  are in  $C\Phi_{tu}$  [4]. Also, it is known that  $(C\Phi_{tu} \cup C)$  is a group under  $\oplus$ . Here we will show the similar relationship as follows: Let  $S$  be the class of all possible cut sets, edge disjoint unions of cut sets in a linear graph  $G$  and  $\emptyset$ . Let  $d$  be a psuedo cut. Now we define  $SD(d)$  as

$$SD(d) = \{x: x = d \oplus s_r, s_r \in S\}$$

Then  $\{SD(d) \cup S\}$  is a group under  $\oplus$ .

The second part shows the relationship among classes of psuedo cuts which leads to an important property that the collection of all possible classes of psuedo cuts of a nonseparable linear graph is an Abelian group under the cross ring sum  $\oplus$  operation. This property is exactly the same as that of the collection of the classes of paths [5].



The last part shows that a psuedo cut is a subset of a set of chords in a linear graph. Then we apply this property to obtain all possible sets of chords in a nonseparable graph.

### PRELIMINARY

Definition 1: Class  $S$  is the collection of all possible cut sets, all possible edge disjoint unions of cut sets and the empty set  $\emptyset$  in a linear graph. Class  $S_s$  is a subclass of  $S$  which consists of all possible cut sets and  $\emptyset$ .

Definition 2: Let  $U$  be a class of sets. Then, "Min  $U$ " is a subclass of  $U$  such that (1) if  $\emptyset$  is in  $U$ ,  $\emptyset$  is in Min  $U$ , (2) for any set  $u_1 \neq \emptyset$  in  $(U - \text{Min } U)^*$ , there exists a set  $u_2 \neq \emptyset$  in Min  $U$  such that  $u_1 \supset u_2$ , and (3) for any two sets  $u_r \neq \emptyset$  and  $u_t \neq \emptyset$  in Min  $U$ ,  $u_r \not\supset u_t$ .

Example 1: Suppose sets  $u_1 = \{abc\}$  and  $u_2 = \{a\}$  are in  $U$ . Then  $u_1$  is not in Min  $U$  because  $u_2 \subset u_1$ .

From the above definitions, it can be seen that

$$S_s = \text{Min } S \quad (1)$$

Definition 3: Let  $U$  and  $V$  be classes of sets. Then  $U \otimes V$  is defined as

$$U \otimes V = \text{Min } \{x: x=u \oplus v, u \in U, v \in V\} \quad (2)$$

---

\*  $A-B$  is a class of sets each of which is in  $A$  but not in  $B$ .

where  $u \oplus v = (u-v) \cup (v-u)$ . For convenience,  $\oplus$  is called a cross ring sum. When there is only one set in each of  $U$  and  $V$ ,  $\oplus$  becomes  $\oplus$ . \*\*

#### PSUEDO CUT IN CLASS $D(e)$

Definition 4: Let  $G$  be a linear graph and edge  $e$  is in  $G$ . Then class  $D(e)$  is defined as

$$D(e) = \text{Min} \{x: x = \{e\} \oplus s, s \in S\} \quad (3)$$

under the condition that set  $\{e\}$  is not a cut set in  $G$ .

$D(e)$  is called a class of psuedo cuts of  $G$  with respect to edge  $e$ , and each set in  $D(e)$  is called a psuedo cut.

Since  $\emptyset$  is in  $S$ , set  $\{e\}$  is in  $D(e)$ . Thus edge  $e$  is a psuedo cut. It is also clear that every psuedo cut in  $D(e)$  except  $\{e\}$  becomes a cut set by adding edge  $e$ .

Example 2: Class  $S_s$  of a linear graph  $G$  in Fig. 1 consists of the following sets:

$$\emptyset, \{e_0 e_1 e_3\}, \{e_0 e_1 e_4\}, \{e_0 e_2 e_3\}, \{e_0 e_2 e_4\}, \{e_1 e_2\} \text{ and } \{e_3 e_4\}$$

Class  $S$  of  $G$  consists of all sets in  $S_s$  and  $\{e_1 e_2 e_3 e_4\}$ .

The class  $D(e_0)$  which can be obtained by Equation (3) or directly from  $G$ , consists of  $\{e_0\}$ ,  $\{e_1 e_3\}$ ,  $\{e_1 e_4\}$ ,  $\{e_2 e_3\}$ , and  $\{e_2 e_4\}$ .

---

\*\* Ring sum  $\oplus$  of  $x$  and  $y$  is defined as  $x \oplus y \subset \overline{xy} \cup \overline{yx} = (x-y) \cup (y-x)$ .




Fig. 1. A Linear Graph G

It is known that the ring sum of two paths both of which are between two vertices  $p$  and  $q$  in a non-oriented linear graph is either a circuit or an edge disjoint union of circuits. Two psuedo cuts with respect to edge  $e$  have the similar property which is given by the following theorem.

Theorem 1: Let  $d_1$  and  $d_2$  be psuedo cuts in  $D(e)$ . Then  $d_1 \oplus d_2$  is a set in  $S$ .

Proof: Since  $d_i \cup \{e\}$  ( $i=1,2$ ) is a cut set of  $G$  by Equation (3), we have

$$d_1 \oplus d_2 = (d_1 \oplus \{e\}) \oplus (d_2 \oplus \{e\}) = s_1 \oplus s_2 \quad (4)$$

Notice if  $d_i \neq \{e\}$ ,  $d_i \oplus e = d_i \cup \{e\}$  which is a cut set  $s_i$  in  $S$ . If  $d_i = \{e\}$ , then  $d_i \oplus \{e\} = \emptyset$  which is also in  $S$ . It is obvious that  $s_1 \oplus s_2$  is in  $S$ . Q.E.D.

Lemma 1: Let  $d$  be a set in  $D(e)$  and  $s$  be a set in  $S$ . Then  $d \oplus s$  is either a psuedo cut in  $D(e)$  or an edge disjoint union of a psuedo cut in  $D(e)$  and a set in  $S$ .

The proof is similar to that of Theorem 1.



In order to obtain a class which contains psuedo cuts and sets, each of which is an edge disjoint union of a psuedo cut and a set in  $S$ , we define the following:

Definition 5: For a given  $G$  and  $S$  of  $G$ , class  $SD(e)$  is defined as

$$SD(e) = \{x: x = \{e\} \oplus s, s \in S\} \quad (5)$$

The difference between  $D(e)$  and  $SD(e)$  is that there is no (Min.) in Equation (5). Thus,  $SD(e)$  is the collection of all possible psuedo cuts in  $D(e)$  and all possible sets each of which is an edge disjoint union of a psuedo cut in  $D(e)$  and cut sets in  $S$ .

From Theorem 1 and Lemma 1, we have the following:

Lemma 2: Let  $SD(e) \cup S$  be the class of all sets in  $SD(e)$  and  $S$  of  $G$ . Then  $SD(e) \cup S$  is an Abelian group under the operation  $\oplus$ .

It can be seen by Lemma 2 that the relationship among psuedo cuts in  $D(e)$  are the same as that of paths between two vertices and circuits in a linear graph.

#### PROPERTIES OF CLASS $D(e)$ OF PSUEDO CUTS

By Definition 1, class  $S_g$  is a subset of class  $S$ . Notice that  $S$  is a class of cut sets, edge disjoint unions of cut sets, and the empty set. In order to obtain class  $D(e)$  (Definition 4) by Equation (3) we must know class  $S$  of a linear graph  $G$ . Because of "Min" in Equation (3), we can replace  $S$  in Equation (3) by  $S_g$  which is given by the following theorem.

Theorem 2: Class  $D(e)$  of psuedo cuts with respect to edge  $e$  can be expressed as

$$D(e) = \text{Min } \{x: x = \{e\} \oplus s, s \in S_s\} \quad (6)$$

Proof: Consider  $\{e\} \oplus s'$  where  $s'$  is an edge disjoint union of cut sets  $s_1, s_2, \dots$ , and  $s_m$ . Then  $\{e\} \oplus s'$  can be expressed as  $\{e\} \oplus s_1 \oplus s_2 \oplus \dots \oplus s_m$ . If  $s'$  does not contain edge  $e$ , then  $d \oplus s'$  is not in  $D(e)$  because  $\{e\}$  is in  $D(e)$ . Suppose edge  $e$  is in  $s'$ . Without loss of generality, let edge  $e$  be in  $s_1$ . Notice that  $s_1$  is in  $S_s$ . Since  $e \oplus s_1 \subset e \oplus s'$ , set  $\{e\} \oplus s'$  is not in  $D(e)$ . Q.E.D.

By Theorem 2, we can use Equation (6) as the definition of  $D(e)$  which is obviously simpler to obtain  $D(e)$ . However, we will see later that  $S$  can not be replaced by  $S_s$  when a class  $D(e_1 e_2 \dots e_m)$  of psuedo cuts with respect to edges  $e_1, e_2, \dots$ , and  $e_m$  for  $m > 1$  is concerned.

There is a relationship between a psuedo cut in  $D(e)$  and class  $D(e)$  as follows.

Lemma 3: Let  $d$  be a psuedo cut in  $D(e)$ . Then  $D(e)$  can be expressed as

$$D(e) = \text{Min } \{x: x = d \oplus s, s \in S\} \quad (7)$$

Proof: From Equation (3), we can see that any psuedo cut  $d$  in  $D(e)$  can be expressed as  $\{e\} \oplus s'$  where  $s' \in S$ . Thus the right hand side of Equation (7) is equal to

$$\text{Min } \{x: x = \{e\} \oplus s' \oplus s, s \in S\} = \text{Min } \{x: x = \{e\} \oplus s'', s'' \in S\} \quad (8)$$

which is the definition of  $D(e)$ . Notice that  $S$  is an Abelian group under  $\oplus$ . Q.E.D.

The use of Theorem 2 gives the relationship between classes  $D(e)$  and  $S_s$  as follows:

Lemma 4:

$$D(e) \otimes S_s = D(e) \quad (9)$$

Proof: This lemma can be proven easily by knowing that (a)  $\emptyset$  is in  $S_s$ , (b)

$$S_s \otimes S_s = S_s \quad (10)$$

and (c)  $S$  in Equation (3) can be substituted by  $S_s$  by Theorem 2.

When  $G$  is nonseparable [2],  $D(e)$  has very interesting properties as

Theorem 3: Let  $G$  be nonseparable. Then

$$D(e) \otimes D(e) = S_s \quad (11)$$

Before proving Theorem 3, we will prove the following lemma.

Lemma 5: If  $s_1$  and  $s_2$  are fundamental cut sets with respect to a tree in  $G$  [2], then  $s_1 \oplus s_2$  is a cut set.

Proof: Let  $s_1$  and  $s_2$  be fundamental cut sets with respect to a tree  $T$ . Because each of  $s_1$  and  $s_2$  contains only one edge in  $T$ , deletion of all edges in  $s_1$  and  $s_2$  from  $G$  produces exactly three connected parts (an isolated vertex will be considered as one connected part). Let these parts be  $g_1, g_2$ , and  $g_3$ . For convenience, symbol



$\mathcal{E}(g_i, g_j)$  represents all edges each of which is connected between a vertex in  $g_i$  and a vertex in  $g_j$ . Without the loss of generality, let

$$s_1 = [\mathcal{E}(g_1, g_2) \mathcal{E}(g_1, g_3)] \quad (12)$$

and

$$s_2 = [\mathcal{E}(g_1, g_2) \mathcal{E}(g_2, g_3)] \quad (13)$$

Then  $s_1 \oplus s_2$  will be

$$s_1 \oplus s_2 = \mathcal{E}(g_1, g_3) \mathcal{E}(g_2, g_3) \quad (14)$$

which is clearly a cut set in  $G$ .

Q.E.D.

Proof of Theorem 3: From Theorem 1, it is clear that the ring sum of any two distinct psuedo cuts in  $D(e)$  is either a cut set or an edge disjoint union of cut sets. Hence, it is only necessary to show that for any cut set in  $G$ , there exists two psuedo cuts in  $D(e)$  such that the ring sum of these two psuedo cuts will give this cut set. Let  $s$  be a cut set.

Case 1: Suppose  $e \in s$ . Then  $d = \{e\} \oplus s$  is in  $D(e)$ . Also  $\{e\}$  is in  $D(e)$ . Hence  $\{e\} \oplus d = s$ .

Case 2: Suppose  $s$  does not contain edge  $e$ . Since  $G$  is non-separable, there exists a circuit  $C$  which contains edges  $e$  and  $e'_t$  where  $e'_t$  is in  $s$ . Let an edge sequence of the circuit  $C$  [2] be

$$e'_1 e'_2 \dots e'_t \dots e'_k, e, e_{k+1} \dots e'_u \dots e'_m$$

where  $e'_t$  and  $e'_u$  are in  $s$  and  $e'_{t+1}, \dots, e'_k, e, e_{k+1}, \dots$ , and  $e'_{u-1}$  are not in  $s$  as shown in Figure 2A.




Fig. 2. Cut set  $s$  and edge  $e$  in  $G$ .

Let  $\zeta_1$  be the endpoint [2] of edges  $e'_{t-1}$  and  $e'_t$  and  $\zeta_2$  be the endpoint of edges  $e'_u$  and  $e'_{u+1}$ . Also let  $g_1$  and  $g_2$  be the two connected parts obtained by deleting all edges in  $s$  where edge  $e$  is in  $g_2$ . Since  $g_1$  is connected, there exists at least one path between  $\zeta_1$  and  $\zeta_2$  as shown in Figure 2B. Hence, there exists a circuit  $c'$  in  $G$  which contains edge  $e$  and exactly two edges in  $s$ . Notice that if  $\zeta_1$  and  $\zeta_2$  are the same, the existence of such a circuit is obvious. By choosing a tree  $T$  of  $G$  which contains all edges in the circuit  $c'$  except edge  $e'_t$ , we can obtain a fundamental cut set  $s'$  which contains edges  $e$  and  $e'_t$ . Notice that  $s$  is also a fundamental cut set with respect to  $T$ . Hence, by Lemma 5,  $s''$ , which is equal to  $s \oplus s'$ , is a cut set in  $G$  and edge  $e$  is in  $s''$ . Thus, there are two psuedo cuts  $\{e\} \oplus s'$  and  $\{e\} \oplus s''$  such that the ring sum of these two psuedo cuts gives the cut set  $s$ . Q.E.D.

Example 3:  $D(e_0)$  in Example 1 is

$$D(e_0) = \{\{e_0\}, \{e_1e_3\}, \{e_1e_4\}, \{e_2e_3\}, \{e_2e_4\}\}$$

Thus

$$D(e_0) \otimes D(e_0) = \{\emptyset, \{e_0e_1e_2\}, \{e_0e_1e_4\}, \{e_0e_2e_3\}, \{e_0e_2e_4\}, \{e_3e_4\}, \{e_1e_2\}\}$$

which is equal to  $S_S$ . A common property of psuedo cuts in  $D(e)$  is given by the following theorem.

Theorem 4: Let  $s \in S$  and  $D(e) \neq S$ . A set  $\{e\} \otimes s$  is in  $D(e)$  if and only if  $\{e\} \otimes s$  does not contain a cut set in  $S$ .

Proof: Suppose  $\{e\} \otimes s$  contains a cut set  $s'$ . Then  $\{e\} \otimes s \otimes s' \subset \{e\} \otimes s$ . Thus  $\{e\} \otimes s$  will not be in  $D(e)$ . Suppose  $\{e\} \otimes s$  is in  $D(e)$ . Then  $s$  is a cut set but is not an edge disjoint union of cut sets by Theorem 2. Furthermore, edge  $e$  is in  $s$ , thus there is no cut set in  $\{e\} \otimes s$ . Q.E.D.

#### PSUEDO CUT WITH RESPECT TO SEVERAL EDGES

The previous section gives the definition and properties of psuedo cuts in  $D(e)$ . Here we will generalize the definition of classes of psuedo cuts to include  $D(e_1e_2\dots e_m)$ .

Definition 6: A class  $D(e_1\dots e_m)$  in  $G$  is defined as

$$D(e_1\dots e_m) = \text{Min} \{x: x = \{e_1\dots e_m\} \otimes s, s \in S\} \quad (15)$$

where edges  $e_1, e_2, \dots$ , and  $e_m$  are in  $G$ . When  $D(e_1\dots e_m) \neq S_p$ , a set in  $D(e_1\dots e_m)$  is called a psuedo cut.  $D(e_1\dots e_m)$  is a class of psuedo cuts with respect to edges  $e_1, e_2, \dots$ , and  $e_m$ .



Theorem 5: Let  $D(e_1 e_2) \neq S_s$ ,  $D(e_1) \neq S_s$  and  $D(e_2) \neq S_s$ .

Then

$$D(e_1 e_2) = D(e_1) \otimes D(e_2) \quad (16)$$

Proof: Any set in  $D(e_1) \otimes D(e_2)$  can be expressed as

$$\{e_1\} \oplus s_1 \oplus \{e_2\} \oplus s_2 = \{e_1 e_2\} \oplus s \text{ where } s_1 \oplus s_2 = s, s_1, s_2 \text{ and } s \in S.$$

Thus

$$D(e_1 e_2) \supset D(e_1) \otimes D(e_2) \quad (17)$$

Suppose set  $d = \{e_1 e_2\} \oplus s$  is in  $D(e_1 e_2)$  but is not in  $D(e_1) \otimes D(e_2)$ .

Let  $s$  be expressed as  $s_1 \oplus s_2$  where  $e_1 \in s_1$ , and  $s_1 \in S_s$ . When  $e_1 \in d$ ,  $s_1$  is  $\emptyset$  by definition. Then  $\{e_1\} \oplus s_1 \in D(e_1)$ . In order that  $d$  is not in  $D(e_1) \otimes D(e_2)$ ,  $\{e_2\} \oplus s_2$  can not be in  $D(e_2)$ . Because  $\{e_2\} \oplus s_2$  is in  $D(e_2)$  there must be a proper non-empty subset of  $\{e_1 e_2\} \oplus s$  in  $D(e_1) \otimes D(e_2)$ . However, if there is such a subset  $d'$  in  $D(e_1) \otimes D(e_2)$ ,  $d' \subset d$ ,  $d$  is not in  $D(e_1 e_2)$  which is a contradiction. Thus  $\{e_2\} \oplus s_2$  can not be in  $D(e_2)$ . Then by Theorem 4,  $\{e_2\} \oplus s_2$  must contain a cut set. Let this cut set be  $s'$ . Then  $\{e_1 e_2\} \oplus s \supset \{e_1 e_2\} \oplus s \oplus s'$ . Hence by Definitions 2 and 3,  $d$  is not in  $D(e_1 e_2)$  (contradiction). Thus

$$D(e_1 e_2) \subset D(e_1) \otimes D(e_2) \quad (18)$$

Q.E.D.

From Theorem 5 we have

Lemma 6: Let  $e_1$  and  $e_2$  be in  $G$ . Then

$$\begin{aligned} & \text{Min } \{ \text{Min } y: y = \{e_1\} \oplus s, s \in S \} \otimes \text{Min } \{ z: z = \{e_2\} \oplus s, s \in S \} \\ & = \text{Min } \{ \{ y: y = \{e_1\} \oplus s, s \in S \} \otimes \{ z: z = \{e_2\} \oplus s, s \in S \} \} \end{aligned} \quad (19)$$

Similar to Theorem 4, we have the following.

Theorem 6:  $\{e_1 e_2\} \oplus s$  is in  $D(e_1 e_2)$  if and only if  $\{e_1 e_2\} \oplus s$  does not contain a cut set in  $S$  where  $s \in S$ .

Proof: Suppose  $\{e_1 e_2\} \oplus s$  contain a cut set  $s'$ . Then  $\{e_1 e_2\} \oplus (s \oplus s') = \{e_1 e_2\} \oplus s' \subset \{e_1 e_2\} \oplus s$ . Hence  $\{e_1 e_2\} \oplus s$  will not be in  $D(e_1 e_2)$ . Suppose  $\{e_1 e_2\} \oplus s$  is in  $D(e_1 e_2)$ , then there is no proper subset (non-empty) of  $\{e_1 e_2\} \oplus s$  in  $D(e_1 e_2)$ . Thus  $\{e_1 e_2\} \oplus s$  can not contain a cut set in  $S$ . Q.E.D.

We will generalize Theorem 6 as follows.

Theorem 7: Let  $s \in S$ . A set  $\{e_1 \dots e_p\} \oplus s$  is in  $D(e_1 \dots e_p)$   $\neq S_s$  if and only if  $\{e_1 \dots e_p\} \oplus s$  does not contain a cut set in  $S$ .

Proof: If  $\{e_1 \dots e_p\} \oplus s$  contains a cut set  $s'$ , then  $\{e_1 \dots e_p\} \oplus s \oplus s' \subset \{e_1 \dots e_p\} \oplus s$ . Thus  $\{e_1 \dots e_p\} \oplus s$  will not be in  $D(e_1 \dots e_p)$ . Suppose  $\{e_1 \dots e_p\} \oplus s$  is in  $D(e_1 \dots e_p)$ . Then there is no non-empty proper subset of  $\{e_1 \dots e_p\} \oplus s$  is in  $D(e_1 \dots e_p)$ . Thus it is impossible for  $\{e_1 \dots e_p\} \oplus s$  to contain a cut set. Q.E.D.

The relationship between  $D(e_1 \dots e_p)$  and  $D(e_1)$ ,  $D(e_2)$ , ..., and  $D(e_p)$  is given by the following.

Theorem 8: Let  $D(e_1 \dots e_p) \neq S_s$ ,  $D(e_1 \dots e_{p-1}) \neq S_s$  and  $D(e_p) \neq S_s$  then

$$D(e_1 \dots e_p) = D(e_1 \dots e_{p-1}) \oplus D(e_p) \quad (20)$$

Proof: It is obvious that any set in  $D(e_1 \dots e_{p-1}) \oplus D(e_p)$  can be expressed as  $\{e_1 \dots e_p\} \oplus s$  where  $s \in S$ . Thus

$$D(e_1 \dots e_p) \supset D(e_1 \dots e_{p-1}) \oplus D(e_p) \quad (21)$$



Suppose set  $d = \{e_1 \dots e_p\} \otimes s \in D(e_1 \dots e_p)$  is not in  $D(e_1 \dots e_{p-1}) \otimes D(e_p)$ . Let  $s$  be

$$s = s_1 \oplus s_2 \oplus \dots \oplus s_n \quad (22)$$

where  $s_1, s_2, \dots, s_n \in S_s$  have no edge in common. Furthermore, let  $e_p \in s_1$  ( $s_1$  can be the empty set if  $e_p \notin s$ ). Notice that  $s_i, i=1,2,\dots,n$  is a cut set if  $s_i$  is not the empty set. Then by Theorem 4,  $\{e_p\} \oplus s_1 \in D(e_p)$ . By Theorem 7,  $\{e_1 \dots e_{p-1}\} \oplus s_2 \oplus \dots \oplus s_n$  is not in  $D(e_1 \dots e_{p-1})$  if  $\{e_1 \dots e_{p-1}\} \oplus s_2 \oplus \dots \oplus s_n$  contains a cut set. Thus, in order that  $d$  is not in  $D(e_1 \dots e_{p-1}) \otimes D(e_p)$ ,  $\{e_1 \dots e_{p-1}\} \oplus s_2 \oplus \dots \oplus s_n$  must contain a cut set  $s'$ . Then  $\{e_1 \dots e_p\} \oplus s \oplus s' \subset d$  is in  $D(e_1 \dots e_p)$  which contradicts the assumption that  $d$  is in  $D(e_1 \dots e_{p-1})$ . Hence,  $\{e_1 \dots e_{p-1}\} \oplus s_2 \oplus \dots \oplus s_n$  is in  $D(e_1 \dots e_{p-1})$ . Thus, in order that  $d$  is in  $D(e_1 \dots e_p)$ ,  $d$  must be in  $D(e_1 \dots e_{p-1}) \otimes D(e_p)$ . Q.E.D.

We can generalize the above theorem as follows.

Theorem 9: Let  $D(e_1 \dots e_p) \neq S_s$ ,  $D(e_1 \dots e_i) \neq S_s$  and  $D(e_{i+1} \dots e_p) \neq S_s$  where  $1 \leq i < p$ . Then

$$D(e_1 \dots e_p) = D(e_1 \dots e_i) \otimes D(e_{i+1} \dots e_p) \quad (23)$$

The proof is similar to that of Theorem 8.

Notice that the classes of psuedo cuts in the Theorems 8 and 9 must not be equal to  $S_s$ . In order to show that classes of psuedo cuts including  $S_s$  form an Abelian group under  $\otimes$ , we must remove such restrictions. It is necessary to study the following properties before discussing such a problem.



Lemma 7: Let  $d$  be in  $D(e_1 \dots e_p)$ , then

$$D(e_1 \dots e_p) = \text{Min} \{x: x=d \oplus s, s \in S\} \quad (24)$$

The proof is obvious from Definition 6.

Lemma 8: If  $\{e_1 \dots e_p\}$  is a cut set or an edge disjoint union of cut sets,

$$D(e_1 \dots e_p) = S_s \quad (25)$$

Lemma 9: If  $\{e_1 \dots e_p\}$  contains a cut set  $\{e_{i+1} \dots e_p\}$  where  $1 \leq i < p$ , then

$$D(e_1 \dots e_p) = D(e_1 \dots e_i) \quad (26)$$

Proof: Since every set in  $D(e_1 \dots e_p)$  can be expressed as

$$\{e_1 \dots e_p\} \oplus s = \{e_1 \dots e_i\} \oplus \{e_{i+1} \dots e_p\} \oplus s = \{e_1 \dots e_i\} \oplus s' \quad (27)$$

where  $\{e_{i+1} \dots e_p\} \oplus s = s'$ , the proof follows from Definition 6.

Lemma 10:

$$D(e_1 \dots e_p) \otimes S_s = D(e_1 \dots e_p) \quad (28)$$

The proof is clear because  $\emptyset \in S_s$ .

Theorem 3 gives that  $D(e) \otimes D(e)$  is equal to  $S_s$ . The following is the generalization of Theorem 3.

Theorem 10: Let  $G$  be nonseparable. Then

$$D(e_1 \dots e_p) \otimes D(e_1 \dots e_p) = S_s \quad (29)$$

Proof: Case 1: When  $D(e_1 \dots e_p) = S_s$ , then by Lemma 10, the left hand side of Equation (29) becomes

$$D(e_1 \dots e_p) \otimes S_s = D(e_1 \dots e_p) \quad (30)$$

which is equal to  $S_s$ . Case 2: When  $D(e_1 \dots e_p) \neq S_s$ , then any set in  $D(e_1 \dots e_p) \otimes D(e_1 \dots e_p)$  can be expressed as

$$\{e_1 \dots e_p\} \oplus s_1 \oplus \{e_1 \dots e_p\} \oplus s_2 = s_1 \oplus s_2 \quad (31)$$

which is in  $S_s$ . Hence

$$D(e_1 \dots e_p) \otimes D(e_1 \dots e_p) \subset S_s \quad (32)$$

Because of Lemma 9, we can assume that  $(e_1 \dots e_p)$  does not contain a cut set in  $S_s$ . Now we will prove that  $S_s$  is a subclass of  $D(e_1 \dots e_p) \otimes D(e_1 \dots e_p)$ . In the proof of Theorem 3, we show that for any set  $s$  in  $S_s$ , either  $e_1$  is in  $s$  or there exists  $s_1$  and  $s_2$  in  $S_s$  such that  $e_1 \in s_1$ ,  $e_1 \in s_2$  and  $s_1 \oplus s_2 = s$ . Hence if  $e_1 \in s$ ,  $\{e_1\} \oplus s \in D(e_1)$  which gives that  $\{e_1 \dots e_p\} \oplus s$  is in  $D(e_1 \dots e_p)$ . Also, we know by Theorem 7 that  $\{e_1 \dots e_p\} \in D(e_1 \dots e_p)$ . Thus,  $s$  is in  $D(e_1 \dots e_p) \otimes D(e_1 \dots e_p)$ . If  $e_1 \notin s$ , then  $\{e_1\} \oplus s_1$  and  $\{e_1\} \oplus s_2$  are in  $D(e_1)$ . Thus,  $\{e_1 \dots e_p\} \oplus s_1$  and  $\{e_1 \dots e_p\} \oplus s_2$  are in  $D(e_1 \dots e_p)$ . Hence  $s$  is in  $D(e_1 \dots e_p) \otimes D(e_1 \dots e_p)$ . Thus

$$D(e_1 \dots e_p) \otimes D(e_1 \dots e_p) \supset S_s \quad (33)$$

Q.E.D.

Notice that  $G$  must be nonseparable. For example, if  $G$  is the graph in Figure 3,  $D(e_1) = \{\{e_1\}, \{e_2\}\}$  and  $D(e_1) \otimes D(e_1) = \{\emptyset, \{e_1 e_2\}\} \neq S_s$ .




Fig. 3. A Separable Graph

Now we are ready to remove the restriction on classes of psuedo cuts in Theorem 9.

Theorem 11: Let  $G$  be nonseparable. Then

$$D(e_1 \dots e_p) = D(e_1 \dots e_i) \otimes D(e_{i+1} \dots e_p) \quad (34)$$

where  $1 \leq i < p$ .

Proof: Suppose  $D(e_1 \dots e_p) = S_s$ . Then if  $D(e_{i+1} \dots e_p) = S_s$ , Equation (34) holds because of Lemmas 9 and 10. If both  $D(e_1 \dots e_i)$  and  $D(e_{i+1} \dots e_p)$  are not equal to  $S_s$ , Theorem 10 will give the proof. Thus, we only need to consider the case when  $D(e_1 \dots e_p) = S_s$ . In this case,  $\{e_1 \dots e_p\}$  is either a cut set or an edge disjoint union of cut sets. When  $D(e_1 \dots e_i) = S_s$ , Equation (34) holds because of Lemmas 9 and 10. Thus, we assume that  $D(e_1 \dots e_p) \neq S_s$  and  $D(e_{i+1} \dots e_p) \neq S_s$ . Also by Lemma 9, we assume that  $\{e_1 \dots e_i\}$  and  $\{e_{i+1} \dots e_p\}$  do not contain a cut set. Since  $\{e_1 \dots e_p\} \in S$ , either



$$\{e_1 \dots e_i\} \oplus \{e_1 \dots e_p\} = \{e_{i+1} \dots e_p\} \quad (35)$$

or a proper nonzero subset of  $\{e_{i+1} \dots e_p\}$  is in  $D(e_1 \dots e_i)$ . However, because  $\{e_{i+1} \dots e_p\}$  does not contain a cut set,  $\{e_{i+1} \dots e_p\}$  must be in  $D(e_1 \dots e_i)$  by Theorem 7. Thus, by Lemma 7,

$$D(e_1 \dots e_i) = D(e_{i+1} \dots e_p) \quad (36)$$

Hence, by Theorem 10, Equation (33) with  $D(e_1 \dots e_p) = S_s$  holds. Q.E.D.

The direct application of Theorem 11 gives the following important theorem.

Theorem 12: Let  $G$  be nonseparable. Then

$$\begin{aligned} [D(e_{11} \dots e_{1m}) \otimes D(e_{21} \dots e_{2m})] \otimes D(e_{31} \dots e_{3p}) = \\ D(e_{11} \dots e_{1m}) \otimes [D(e_{21} \dots e_n) \otimes D(e_{31} \dots e_{3p})] \end{aligned} \quad (37)$$

Definition 7: Let  $\mathcal{E}$  be a set of edges  $e_1, e_2, \dots$ , and  $e_p$  in  $G$ . Then  $D(\mathcal{E})$  is a short hand notation of  $D(e_1 e_2 \dots e_p)$ . Let  $\mathcal{E} = \mathcal{E}_1 \oplus \mathcal{E}_2$ . Then  $D(\mathcal{E}_1 \oplus \mathcal{E}_2) = D(\mathcal{E})$  where  $\oplus$  is any operation defined in the set theory. With this notation, we have

Lemma 11: Let  $G$  be nonseparable. Then

$$D(\mathcal{E}_1) \otimes D(\mathcal{E}_2) = D(\mathcal{E}_1 \cup \mathcal{E}_2) = D(\mathcal{E}_1 \oplus \mathcal{E}_2) \quad (38)$$

The proof can be obtained by the use of Lemma 9 and Theorem 11. From Theorems 11 and 12, we know that

$$D(\mathcal{E}_1) \otimes D(\mathcal{E}_2) \otimes D(\mathcal{E}_3) = D(\mathcal{E}_1 \cup \mathcal{E}_2 \cup \mathcal{E}_3) \quad (39)$$

where  $\mathcal{E}_1$ ,  $\mathcal{E}_2$ , and  $\mathcal{E}_3$  are sets of edges in a nonseparable graph  $G$ .

Hence, with Definition 6, we have the following theorem.

Theorem 13: Let  $G$  be nonseparable. Then

$$\begin{aligned} \text{Min} \{ \text{Min} \{x: x = \mathcal{E}_1 \oplus s, s \in S\} \oplus \text{Min} \{y: y = \mathcal{E}_2 \oplus s, s \in S\} \oplus \\ \text{Min} \{z: z = \mathcal{E}_3 \oplus s, s \in S\} \} = \text{Min} \{ \{x: x = \mathcal{E}_1 \oplus s, s \in S\} \oplus \\ \{y: y = \mathcal{E}_2 \oplus s, s \in S\} \oplus \{z: z = \mathcal{E}_3 \oplus s, s \in S\} \} \end{aligned} \quad (40)$$

where  $\mathcal{E}_1$ ,  $\mathcal{E}_2$ , and  $\mathcal{E}_3$  are sets of edges in  $G$ .

Definition 8: Collection  $\{D\}$  is defined as the collection of all distinct classes of psuedo cuts in  $G$  and  $S_s$ . With this definition, it is easily seen that the previous results are enough to state that,

Theorem 14: Let  $G$  be nonseparable, then  $\{D\}$  of  $G$  is an Abelian group under the operation  $\oplus$ , where  $S_s$  is a unit element and the inverse of  $D(e_1 \dots e_p)$  is itself.

Example 4:  $S_s$  of  $G$  in Figure 4 is

$$\begin{aligned} S_s = S = \{ \emptyset, \{e_1 e_2 e_3\}, \{e_1 e_4 e_6\}, \{e_1 e_2 e_5 e_6\}, \{e_2 e_3 e_4 e_6\}, \\ \{e_3 e_5 e_6\}, \{e_2 e_4 e_5\}, \{e_1 e_3 e_4 e_5\} \} \end{aligned}$$

Then  $D(e_1)$  is

$$D(e_1) = \text{Min} \{x: x = \{e_1\} \oplus s, s \in S_s\} = \{ \{e_1\}, \{e_2 e_3\}, \{e_4 e_6\}, \{e_2 e_5 e_6\}, \{e_3 e_4 e_5\} \}$$

and  $D(e_2)$  is

$$D(e_2) = \text{Min}\{x: x = \{e_2\} \oplus s, s \in S_s\} = \{\{e_2\}, \{e_1 e_3\}, \{e_1 e_5 e_6\}, \{e_3 e_4 e_6\}, \{e_4 e_5\}\}$$

Thus, by Theorem 5 and Definition (3),  $D(e_1 e_2)$  is equal to

$$\text{Min}\{x: x = d_1 \oplus d_2, d_1 \in D(e_1), d_2 \in D(e_2)\} = D(e_1 e_2) =$$

$$D(e_1) \otimes D(e_2) = \{\{e_1 e_2\}, \{e_3\}, \{e_5 e_6\}, \{e_2 e_4 e_6\}, \{e_1 e_4 e_5\}\}$$

On the other hand,  $D(e_1 e_2)$  is given by Definition 6 as

$$D(e_1 e_2) = \text{Min}\{x: x = \{e_1 e_2\} \oplus s, s \in S\} = \{\{e_1 e_2\}, \{e_3\}, \{e_2 e_4 e_6\}, \{e_5 e_6\}, \{e_1 e_4 e_5\}\}$$

which is equal to  $D(e_1) \otimes D(e_2)$ . Notice that set  $\{e_3\}$  is in  $D(e_1 e_2)$ .

Thus, by Lemma 7,  $D(e_1 e_2)$  can be expressed as

$$D(e_1 e_2) = \text{Min}\{x: x = \{e_3\} \oplus s, s \in S\} = \{\{e_3\}, \{e_1 e_2\}, \{e_2 e_4 e_6\}, \{e_5 e_6\}, \{e_1 e_4 e_5\}\}$$

which is, by Definition 6, equal to  $D(e_3)$ .




Fig. 4. A Linear Graph G



$[D(e_1) \otimes D(e_2)] \otimes D(e_3)$  is equal to  $D(e_1 e_2) \otimes D(e_3) = D(e_3) \otimes D(e_3) = S_s$ . On the other hand,  $D(e_1) \otimes [D(e_2) \otimes D(e_3)]$  is equal to  $D(e_1) \otimes D(e_2 e_3)$  where

$$D(e_2 e_3) = \text{Min}\{x: x = \{e_2 e_3\} \oplus s, s \in S\} = \{\{e_2 e_3\}, \{e_1\}, \{e_4 e_6\}, \{e_2 e_5 e_6\}, \{e_3 e_4 e_5\}\}$$

Since  $\{e_1\}$  is in  $D(e_2 e_3)$ , by Lemma 7,  $D(e_2 e_3)$  is equal to  $D(e_1)$ . Thus,

$$D(e_1) \otimes D(e_2 e_3) = D(e_1) \otimes D(e_1) = S_s.$$

Class  $D(e_4)$  is given by

$$D(e_4) = \text{Min}\{x: x = \{e_4\} \oplus s, s \in S\} = \{\{e_4\}, \{e_1 e_6\}, \{e_2 e_3 e_6\}, \{e_2 e_5\}, \{e_1 e_3 e_5\}\}$$

Consider  $[D(e_1 e_2) \otimes D(e_4)] \otimes D(e_3)$  where  $D(e_1 e_2) \otimes D(e_4)$  is

$$D(e_1 e_2) \otimes D(e_4) = D(e_1 e_2 e_4) = \{\{e_1 e_2 e_4\}, \{e_3 e_4\}, \{e_2 e_6\}, \{e_4 e_5 e_6\}, \{e_1 e_3 e_6\}, \\ \{e_1 e_5\}, \{e_2 e_3 e_5\}\}$$

$D(e_1 e_2 e_4) \otimes D(e_3)$  will be

$$D(e_1 e_2 e_4) \otimes D(e_3) = \text{Min}\{x: x = d_1 \oplus d_2, d_1 \in D(e_1 e_2 e_4), \\ d_2 \in D(e_3)\} = \text{Min}\{\{e_1 e_2 e_3 e_4\}, \{e_4\}, \{e_2 e_3 e_6\}, \{e_3 e_4 e_5 e_6\}, \\ \{e_1 e_6\}, \{e_1 e_3 e_5\}, \{e_2 e_5\}, \{e_1 e_2 e_4 e_5 e_6\}\} = \{\{e_4\}, \{e_2 e_3 e_6\}, \\ \{e_1 e_6\}, \{e_1 e_3 e_5\}, \{e_2 e_5\}\}$$

which is equal to  $D(e_4)$ . On the other hand

$$D(e_4) \otimes D(e_3) = \text{Min}\{x: x = d_1 \oplus d_2, d_1 \in D(e_4), d_2 \in D(e_3)\} = \\ \{\{e_3 e_4\}, \{e_1 e_2 e_4\}, \{e_1 e_3 e_6\}, \{e_2 e_6\}, \{e_4 e_5 e_6\}, \{e_2 e_3 e_5\}, \{e_1 e_5\}\} = D(e_3 e_4)$$

Hence  $D(e_1 e_2) \otimes [D(e_4) \otimes D(e_3)]$  will be

$$D(e_1 e_2) \otimes D(e_3 e_4) = \text{Min}\{x: x = d_1 \oplus d_2, d_1 \in D(e_1 e_2), d_2 \in D(e_3 e_4)\} = \\ \{\{e_4\}, \{e_2 e_3 e_6\}, \{e_1 e_6\}, \{e_1 e_3 e_5\}, \{e_2 e_5\}\}$$

which is equal to  $D(e_4)$ . Thus

$$[D(e_1 e_2) \otimes D(e_4)] \otimes D(e_3) = D(e_1 e_2) \otimes [D(e_4) \otimes D(e_3)].$$

#### PROPERTIES OF PSUEDO CUTS AND THEIR APPLICATION

Consider a connected linear graph  $G$  which consists of  $n_v$  vertices and  $n_e$  edges. Then the number of edges in a cut set can not exceed  $n_e - n_v + 2$  because that any cut set can be a fundamental cut set of some tree in  $G$  and any fundamental cut set can not have more than the number of chords plus one edge. We will show here that a psuedo cut can not contain more than  $n_e - n_v + 1$  edges and any psuedo cut is a subset of a set of chords in  $G$ .

From Definition 6 and Lemma 7, we know that if  $D(e_{i_1} \dots e_{i_m})$  and  $D(e_{j_1} \dots e_{j_n})$  have a psuedo cut in common then,

$$D(e_{i_1} \dots e_{i_m}) = D(e_{j_1} \dots e_{j_n}) \quad (41)$$

Conversely, we have the following lemma.

Lemma 12: Suppose set  $\mathcal{E}$  of edges does not contain a cut set.

Then for any subsets  $\mathcal{E}_1$  and  $\mathcal{E}_j$  in  $\mathcal{E}$  where  $\mathcal{E}_1 \neq \mathcal{E}_2$ ,

$$D(\mathcal{E}_1) \neq D(\mathcal{E}_2) \quad (42)$$

Proof: Suppose it is not true. Then  $\mathcal{E}_1$  and  $\mathcal{E}_2$  are in  $D(\mathcal{E}_1)$ . Thus, it is clear by Definition 6,  $\mathcal{E}_1 \oplus \mathcal{E}_2$  is in  $S$  which contradicts the assumption that  $\mathcal{G}$  does not contain a cut set. Q.E.D.

The maximum number of edges which a psuedo cut can have is given by the following theorem.

Theorem 15: The number of edges in a psuedo cut of a connected linear graph  $G$  can not exceed  $n_e - n_v + 1$  where  $n_e$  is the number of edges and  $n_v$  is the number of vertices in  $G$ . Furthermore, there exists a psuedo cut which contains exactly  $n_e - n_v + 1$  edges.

Proof: Suppose  $\{e_1 \dots e_m\}$  is a psuedo cut in  $D(e_{i_1} \dots e_{i_p})$ . Then by Lemma 7,

$$D(e_{i_1} \dots e_{i_p}) = \text{Min}\{x: x = \{e_1 \dots e_m\} \oplus s, s \in S\} = D(e_1 \dots e_m) \quad (43)$$

Also, by Lemma 10,  $\{e_1 \dots e_m\}$  contains no cut set in  $S$ . Thus,  $m \leq n_e - n_v + 1$ .

Let  $\{e_1 \dots e_c\}$  be a set of chords where  $c = n_e - n_v + 1$ . Then  $\{e_1 \dots e_c\}$  contains no cut sets. Thus,  $\{e_1 \dots e_c\}$  is a psuedo cut in  $D(1_1 \dots 1_c)$  by Lemma 10.

With these lemmas and theorem, we can see that (1) any set of chords in  $G$  belongs to exactly one class of psuedo cuts, and (2) any psuedo cut is a subset of a set of chords.

Definition 9:  $\{D\{e_1 \dots e_q\}\}$  is the subclass of  $\{D\}$  of a linear graph  $G$  such that



$$\{D\{e_1 \dots e_q\}\} = \{D(e_{i_1} \dots e_{i_p}) : \{e_{i_1} \dots e_{i_p}\} \subset \{e_1 \dots e_q\}, p=1,2,\dots,q\} \quad (44)$$

Notice that there are  $2^q - 1$  classes in  $\{D\{e_1 \dots e_q\}\}$ . By Lemma 12, all classes in  $\{D\{e_1 \dots e_q\}\}$  are distinct if  $\{e_1 \dots e_q\}$  is a subset of a set of chords with respect to a tree in  $G$ . Suppose  $\{e_1 \dots e_q\}$  does not contain a cut set. Also suppose  $G$  is connected. Consider a subgraph  $g$  of  $G$  which is obtained by deleting all edges in  $\{e_1 \dots e_q\}$  from  $G$ . Since  $\{e_1 \dots e_q\}$  does not contain a cut set,  $g$  is connected and contains all vertices of  $G$ . Thus, there exists a tree in  $g$  which is also a tree in  $G$ . Thus  $\{e_1 \dots e_q\}$  is a subset of a set of chords with respect to a tree in  $G$ . Hence, we have the following Theorem.

Theorem 16: Let  $G$  be connected. Then there are  $2^q - 1$  distinct classes in  $\{D\{e_1 \dots e_q\}\}$  if and only if  $\{e_1 \dots e_q\}$  is a subset of a set of chords with respect to a tree in  $G$ .

The following theorem is one of the important theorems in order to obtain  $\{D\}$ .

Theorem 17: Let  $\{e_1, \dots, e_q\}$  be a set of chords in a nonseparable graph  $G$ . Then, any class of psuedo cuts of  $G$  is equal to one of the classes in  $\{D\{e_1 \dots e_q\}\}$ .

Proof: Let  $T$  be a tree corresponding to set  $\{e_1 \dots e_q\}$  of chords. Consider a class  $D(e_{j_1} \dots e_{j_u})$ . It is obvious that when  $\{e_{j_1} \dots e_{j_u}\} \subset \{e_1 \dots e_q\}$ ,  $D(e_{j_1} \dots e_{j_u}) \in \{D\{e_1 \dots e_q\}\}$ . Thus, we assume that  $\{e_{j_1} \dots e_{j_u}\} \not\subset \{e_1 \dots e_q\}$ . Because of Lemma 8, we can further assume that  $\{e_{j_1} \dots e_{j_u}\}$  does not contain a cut set. Without the loss of

generality, let  $\{e_{j_1} \dots e_{j_u}\} - \{e_1 \dots e_q\} = \{e_{j_1} \dots e_{j_m}\}$  where  $1 < m \leq u$ .

Notice that edges  $e_{j_1} \dots e_{j_m}$  are in  $T$ . Also, let  $s_1, \dots, s_m$  be funda-

mental cut sets of  $e_{j_1}, e_{j_2}, \dots, e_{j_m}$  with respect to  $T$  such that  $e_{j_p} \in$

$s_p$  for  $p=1, 2, \dots, m$ . Then, by Theorem 11,  $D(e_{j_1} \dots e_{j_u})$  can be expressed as

$$D(e_{j_1} \dots e_{j_m}) = D(e_{j_1}) \otimes D(e_{j_2}) \otimes \dots \otimes D(e_{j_m}) \otimes D(e_{j_{m+1}} \dots e_{j_u}) \quad (45)$$

Furthermore, by Theorem 11

$$D(e_{j_p}) = D(s_p - \{e_{j_p}\}) \quad (46)$$

we have

$$\begin{aligned} D(e_{j_1} \dots e_{j_m}) &= D(s_1 - \{e_{j_1}\}) \otimes D(s_2 - \{e_{j_2}\}) \otimes \dots \otimes \\ &D(s_m - \{e_{j_m}\}) \otimes D(e_{j_{m+1}} \dots e_{j_u}) = D(s_1 - \{e_{j_1}\}) \otimes \\ &\{s_2 - \{e_{j_2}\}\} \otimes \dots \otimes \{s_m - \{e_{j_m}\} \otimes \{e_{j_{m+1}} \dots e_{j_u}\}\} \end{aligned} \quad (47)$$

Notice that every edge in  $s_p - \{e_{j_p}\}$  is in the set  $\{e_1 \dots e_q\}$  of chords with respect to  $T$ . Hence

$$\{s_1 - \{e_{j_1}\}\} \otimes \dots \otimes \{s_m - \{e_{j_m}\}\} \otimes \{e_{j_{m+1}} \dots e_{j_u}\} \subset \{e_1 \dots e_q\} \quad (48)$$

Thus  $D(e_{j_1} \dots e_{j_m}) \in \{D\{e_1 \dots e_q\}\}$

Q.E.D.

This theorem gives that  $\{D\} = \{D(e_1 \dots e_q)\}, S_S\}$  if  $\{e_1 \dots e_q\}$  is a set of chords in a nonseparable graph  $G$ . Thus, in order to obtain all possible distinct classes of psuedo cuts, we only need to obtain  $\{D(e_1 \dots e_q)\}$  where  $\{e_1 \dots e_q\}$  is a set of chords.

There are so-called topological formulas which analyze a linear electrical network which requires knowing all possible sets of chords in  $G$  corresponding to the electrical network. The properties of psuedo cuts discussed here yield to obtain all possible such sets by knowing one set of chords and  $S$  of  $G$ . Also, by Lemma 7, we know that any set of chords is in exactly one class of psuedo cut in  $\{D(e_1 \dots e_q)\}$  where  $\{e_1 \dots e_q\}$  is a set of chords. Hence, it is not necessary to test for duplications of sets of chords. The following example will clarify this point.

Example 5: All possible sets of chords in  $G$  shown in Figure 5 can be obtained by the following procedure. Take a tree, say  $\{acd\}$ . Then we can obtain a set of fundamental cut sets with respect to  $\{acd\}$  as  $\{abf\}$ ,  $\{bce\}$ , and  $\{def\}$ . Then

$$S = \{\emptyset, \{abf\}, \{bce\}, \{def\}, \{acef\}, \{abde\}, \{bcd f\}, \{acd\}\}$$




Fig. 5



The set of chords corresponding to  $\{acd\}$  is  $\{bed\}$ . Thus  $\{D\{bed\}\}$  is the collection of all distinct classes of psuedo cuts which consists of the following.

$$\begin{aligned} D(b) &= \{\{b\}, \{af\}, \{ce\}, \{ade\}, \{cdf\}\} \\ D(e) &= \{\{e\}, \{bc\}, \{df\}, \{acf\}, \{abd\}\} \\ D(f) &= \{\{f\}, \{ab\}, \{de\}, \{ace\}, \{bcd\}\} \\ D(be) &= \{\{be\}, \{c\}, \{bdf\}, \{ad\}, \{aef\}\} \\ D(bf) &= \{\{bf\}, \{a\}, \{bde\}, \{cd\}, \{cef\}\} \\ D(ef) &= \{\{ef\}, \{abe\}, \{bcf\}, \{d\}, \{ac\}\} \end{aligned}$$

and

$$D(bef) = \{\{bef\}, \{ab\}, \{cf\}, \{bd\}, \{adf\}\}$$

Thus, all possible sets of chords are these psuedo cuts which consist of  $n_e - n_v + 1 = 3$ , that is,

$$\begin{aligned} \{ade\} \text{ and } \{cdf\} &\in D(b) \\ \{acb\} \text{ and } \{abd\} &\in D(e) \\ \{ace\} \text{ and } \{bcd\} &\in D(f) \\ \{bdf\} \text{ and } \{aef\} &\in D(be) \\ \{bde\} \text{ and } \{cef\} &\in D(bf) \\ \{abe\} \text{ and } \{bef\} &\in D(ef) \\ \{bef\} \text{ and } \{adf\} &\in D(bef) \end{aligned}$$

are all possible sets of chords in  $G$ .

## REFERENCES

1. Whitney, H., "Planar Graphs," Fundamenta Mathematicae, Vol. 21, pp. 73-84; 1933.
2. Seshu, S. and Reed, M. B., Linear Graphs and Electrical Networks, Addison-Wesley; 1961.
3. Ore, O., "Theory of Graphs," Amer. Math. Soc. Colloquium Publ., Vol. 38; 1962.
4. Mayeda, W., "Synthesis of Switching Functions by Linear Graph Theory," IBM Journal, Vol. 4, No. 3, pp. 321-328; July 1960.
5. Mayeda, W., "Properties of Classes of Paths," CSL Report No. R-212, University of Illinois, Urbana, Illinois; May 1964.

# Distribution list as of February 1, 1964

1	Director Air University Library Maxwell Air Force Base, Alabama Attn: CR-4803a	1	The RAND Corporation 1700 Main Street Santa Monica, California Attn: Library	1	Chief of Naval Operations (Code OP-ONT) Department of the Navy Washington, D. C. 20350
1	Redstone Scientific Information Center U.S. Army Missile Command Redstone Arsenal, Alabama	1	Stanford Electronics Laboratories (Unclassified) Stanford University Stanford, California Attn: SEL Documents Librarian	1	Commanding Officer U. S. Army Personnel Research Office Washington 25, D. C.
1	Electronics Research Laboratory (Unclassified) University of California Berkeley 4, California	1	Dr. L. F. Carter Chief Scientist Air Force Room 4E-324, Pentagon Washington 25, D. C.	1	Commanding Officer & Director Code 142 Library David W. Taylor Model Basin Washington, D. C. 20007
2	Hughes Aircraft Company Florence and Teale Culver City, California Attn: N. E. Devereux Technical Document Center	1	Mr. Robert L. Feik Associate Director for Research Research and Technology Division AFSC Bolling Air Force Base 25, D. C.	1	Chief, Bureau of Ships (Code 686) Department of the Navy Washington, D. C. 20360
3	Autonetics (Unclassified) 9150 East Imperial Highway Downey, California Attn: Tech. Library, 3041-11	1	Captain Paul Johnson (USN-Ret) National Aeronautics and Space Administration 1520 H Street, N. W. Washington 25, D. C.	1	Chief, Bureau of Ships (Code 732) Department of the Navy Washington, D. C. 20360
1	Dr. Arnold T. Nordsieck General Motors Corporation Defense Research Laboratories 6767 Hollister Avenue Goleta, California	1	Major Edwin M. Myers Headquarters USAF (AFRDR) Washington 25, D. C.	1	Chief, Bureau of Naval Weapons Technical Library, DLI-3 Department of the Navy Washington, D. C. 20360
1	University of California (Unclassified) Lawrence Radiation Laboratory P. O. Box 808 Livermore, California	1	Dr. James Ward Office of Deputy Director (Research and Info) Department of Defense Washington 25, D. C.	1	Director, (Code 5140) U. S. Naval Research Laboratory Washington, D. C. 20390
1	Mr. Thomas L. Hartwick Aerospace Corporation P. O. Box 95085 Los Angeles 45, California	1	Dr. Alan T. Waterman Director, National Science Foundation Washington 25, D. C.	1	Chief of Naval Research (Code 437) Department of the Navy Washington, D. C. 20360
1	Lt. Colonel Willard Levin Aerospace Corporation P. O. Box 95085 Los Angeles 45, California	1	Mr. G. D. Watson Defense Research Member Canadian Joint Staff 2450 Massachusetts Ave., N. W. Washington 8, D. C.	1	Dr. H. Wallace Sinaiko (Unclassified) Institute for Defense Analyses Research & Engineering Support Division 1666 Connecticut Ave., N. W. Washington 9, D. C.
1	Professor Zorab Kaprelian University of Southern California University Park Los Angeles 7, California	1	Mr. Arthur G. Wimer Chief Scientist Air Force Systems Command Andrews Air Force Base Washington 25, D. C.	1	Data Processing Systems Division National Bureau of Standards Conn. at Van Ness Room 239, Bldg. 10 Washington 25, D. C. Attn: A. K. Smilow
1	Sylvania Electronic Systems - West Electronic Defense Laboratories P. O. Box 205 Mountain View, California Attn: Documents Center	1	Director, Advanced Research Projects Agency Washington 25, D. C.	1	National Bureau of Standards (Unclassified) Research Information Center & Advisory Service on Information Processing Data Processing Systems Division Washington 25, D. C.
1	Varian Associates 611 Hansen Way Palo Alto, California Attn: Dr. Ira Weissman	1	Air Force Office of Scientific Branch Directorate of Engineering Sciences Washington 25, D. C. Attn: Electronics Division	1	Exchange and Gift Division (Unclassified) The Library of Congress Washington 25, D. C.
1	Huston Denslow (Unclassified) Library Supervisor Jet Propulsion Laboratory California Institute of Technology Pasadena, California	1	Director of Science and Technology Headquarters, USAF Washington 25, D. C. Attn: AFRST-EL/GU	1	NASA Headquarters Office of Applications 400 Maryland Avenue, S. W. Washington 25, D. C. Attn: Mr. A. M. Greg Andrus Code FC
1	Professor Nicholas George California Institute of Technology Electrical Engineering Department Pasadena, California	1	AFRST - SC Headquarters, USAF Washington 25, D. C.	1	APGC (PGAPI) Eglin Air Force Base Florida
1	Space Technology Labs., Inc. One Space Park Redondo Beach, California Attn: Acquisitions Group STL Technical Library	1	Headquarters, R & T Division (Unclassified) Bolling Air Force Base Washington 25, D. C. Attn: RTHR	1	Martin Company P. O. Box 5837 Orlando, Florida Attn: Engineering Library MP-30
2	Commanding Officer and Director U.S. Naval Electronics Laboratory San Diego 52, California Attn: Code 2800, C. S. Manning	1	Headquarters, U. S. Army Material Command Research Division, R & D Directorate Washington 25, D. C. Attn: Physics & Electronics Branch Electronics Section	1	Commanding Officer Office of Naval Research, Chicago Branch 6th Floor, 230 North Michigan Chicago 1, Illinois
1	Commanding Officer and Director U.S. Navy Electronics Laboratory San Diego 52, California Attn: Library	1	Commanding Officer Diamond Ordnance Fuze Laboratories Washington 25, D. C. Attn: Librarian, Room 211, Bldg. 92	1	Laboratories for Applied Sciences University of Chicago 6220 South Drexel Chicago 37, Illinois
1	Office of Naval Research Branch Office 1000 Geary Street San Francisco, California	1	Operation Evaluation Group Chief of Naval Operations (OP-03EG) Department of Navy Washington, D. C. 20350	1	Librarian (Unclassified) School of Electrical Engineering Purdue University Lafayette, Indiana



1	Donald L. Epley (Unclassified) Department of Electrical Engineering State University of Iowa Iowa City, Iowa	1	AFMDC (MDSGP/Capt. Wright) Holloman Air Force Base New Mexico	1	H. E. Cochran Oak Ridge National Laboratory P. O. Box X Oak Ridge, Tennessee
1	Commanding Officer U. S. Army Medical Research Laboratory Fort Knox, Kentucky	1	Commanding General White Sands Missile Range New Mexico	1	U.S. Atomic Energy Commission Office of Technical Information Extension P. O. Box 62 Oak Ridge, Tennessee
2	Keats A. Pullen, Jr. Ballistic Research Laboratories Aberdeen Proving Ground, Maryland	1	Microwave Research Institute Polytechnic Institute of Brooklyn 55 John Street (Unclassified) Brooklyn 1, New York	1	President U.S. Army Air Defense Board Fort Bliss, Texas
1	Director U. S. Army Human Engineering Laboratories Aberdeen Proving Ground, Maryland	1	Cornell Aeronautical Laboratory, Inc. 4455 Genesee Street Buffalo 21, New York Attn: J. P. Desmond, Librarian	1	U.S. Air Force Security Service San Antonio, Texas Attn: ODC-R
1	Mr. James Tippet National Security Agency Fort Meade, Maryland	1	Sperry Gyroscope Company (Unclassified) Marine Division Library 155 Glen Cove Road Glen Cove, L. I., New York Attn: Mrs. Barbara Judd	1	Director Human Resources Research Office The George Washington University 300 North Washington Street Alexandria, Virginia
1	Commander Air Force Cambridge Research Laboratories Laurence G. Hanscom Field Bedford, Massachusetts	1	Major William Harris RADC (RAWI) Griffiss Air Force Base New York	20	Defense Documentation Center Cameron Station Alexandria, Virginia 22314
1	Dr. Lloyd Hollingsworth Director, ERD AFRL L. G. Hanscom Field Bedford, Massachusetts	1	Rome Air Development Center Griffiss Air Force Base New York Attn: Documents Library RAALD	1	Commander U. S. Army Research Office Highland Building 3045 Columbia Pike Arlington 4, Virginia
1	Data Sciences Laboratory Air Force Cambridge Research Lab Office of Aerospace Research, USAF L. G. Hanscom Field Bedford, Massachusetts Attn: Lt. Stephen J. Kahne - CRB	1	Library (Unclassified) Light Military Electronics Department General Electric Company Armament & Control Products Section Johnson City, New York	1	U.S. Naval Weapons Laboratory Computation and Analysis Laboratory Dahlgren, Virginia Attn: Mr. Ralph A. Niemann
1	Instrumentation Laboratory (Unclassified) Massachusetts Institute of Technology 68 Albany Street Cambridge 39, Massachusetts	1	Columbia Radiation Laboratory Columbia University (Unclassified) 538 West 120th Street New York 57, New York	2	Army Material Command Research Division R & D Directorate Bldg. T-7 Gravelley Point, Virginia
1	Research Laboratory of Electronics (Unclassified) Massachusetts Institute of Technology Cambridge 39, Massachusetts Attn: Document Room 26-327	1	Mr. Alan Barnum Rome Air Development Center Griffiss Air Force Base Rome, New York		
1	Dr. Robert Kingston Lincoln Laboratories Lexington, Massachusetts	1	Dr. E. Howard Holt (Unclassified) Director Plasma Research Laboratory Rensselaer Polytechnic Institute Troy, New York		
1	Lincoln Laboratory (Unclassified) Massachusetts Institute of Technology P. O. Box 73 Lexington 73, Massachusetts Attn: Library, A-082	3	Commanding Officer U.S. Army Research Office (Durham) Box CM, Duke Station Durham, North Carolina Attn: CRD-AA-1P, Mr. Ulsh		
1	Sylvania Electric Products, Inc. Electronic Systems Waltham Labs. Library 100 First Avenue Waltham 54, Massachusetts	1	Battelle-DEFENDER Battelle Memorial Institute 505 King Avenue Columbus 1, Ohio		
1	Minneapolis-Honeywell Regulator Co. Aeronautical Division (Unclassified) 2600 Ridgeway Road Minneapolis 13, Minnesota Attn: Dr. D. F. Elwell Main Station: 625	1	Aeronautical Systems Division Navigation and Guidance Laboratory Wright-Patterson Air Force Base Ohio		
1	Inspector of Naval Material Bureau of Ships Technical Representative 1902 West Minnehaha Avenue St. Paul 4, Minnesota	1	Aeronautical Systems Division Directorate of Systems Dynamic Analysis Wright-Patterson Air Force Base Ohio		
20	Activity Supply Officer, USAELRDL Building 2504, Charles Wood Area Fort Monmouth, New Jersey For: Accountable Property Officer Marked: For Inst. for Exploratory Research Inspect at Destination Order No. 576-PM-63-91	1	Commander Research & Technology Div. Wright-Patterson Air Force Base Ohio 45433 Attn: MAYT (Mr. Evans)		
1	Commanding General U. S. Army Electronic Command Fort Monmouth, New Jersey Attn: AMSEL-RE	1	Commanding Officer (AD-5) U.S. Naval Air Development Center Johnsville, Pennsylvania Attn: NADC Library		
1	Mr. A. A. Lundstrom Bell Telephone Laboratories Room 2E-127 Whippany Road Whippany, New Jersey	2	Commanding Officer Frankford Arsenal Philadelphia 37, Pennsylvania Attn: SMUFA-1300		